



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA



NATIONAL SENIOR CERTIFICATE

GRADE 12

MATHEMATICS P2

NOVEMBER 2021

MARKS: 150

TIME: 3 hours

This question paper consists of 13 pages and 1 information sheet.



INSTRUCTIONS AND INFORMATION

Read the following instructions carefully before answering the questions.

1. This question paper consists of 11 questions.
2. Answer ALL the questions in the SPECIAL ANSWER BOOK provided.
3. Clearly show ALL calculations, diagrams, graphs, etc. that you have used in determining your answers.
4. Answers only will NOT necessarily be awarded full marks.
5. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
6. If necessary, round off answers correct to TWO decimal places, unless stated otherwise.
7. Diagrams are NOT necessarily drawn to scale.
8. An information sheet with formulae is included at the end of the question paper.
9. Write neatly and legibly.



QUESTION 1

A bakery kept a record of the number of loaves of bread a tuck-shop ordered daily over the last 18 days. The information is shown in the table below.


10	11	13	14	14	15	16	18	18
19	19	20	21	35	35	37	40	41

- 1.1 Calculate the:


- 1.1.1 Mean number of loaves of bread ordered daily (2)
- 1.1.2 Standard deviation of the data (1)
- 1.1.3 Number of days on which the number of loaves of bread ordered was more than one standard deviation above the mean (2)

- 1.2 The tuck-shop owner was not able to sell all the loaves of bread delivered daily. He calculated the mean number of loaves sold over the 18 days to be 20. Calculate the number of loaves of bread which were NOT sold over the 18 days. (2)
- 1.3 One of the two box and whisker diagrams drawn below represents the data given in the table above.

Graph A:



Graph B:



- 1.3.1 Which ONE of the two box and whisker diagrams, drawn above, correctly represents the data? Write down a reason for your answer. (2)
- 1.3.2 Describe the skewness of the data. (1)
- [10]



QUESTION 2

A farm stall sells milk in 5-litre containers to the local community. The price varies according to the availability of milk at the farm stall. The price of milk, in rands per 5-litre container, and the number of 5-litre containers of milk sold, are recorded in the table below.


Price of milk in rands per 5-litre container (x)	26	32	36	28	40	33	29	34	27	30
Number of 5-litre containers of milk sold (y)	48	30	26	44	23	32	39	29	42	33

- 2.1 On the grid provided in the ANSWER BOOK, draw the scatter plot to represent the data. (3)
- 2.2 Determine the equation of the least squares regression line for the data. (3)
- 2.3 If the farmer sells a 5-litre container of milk for R38, predict the number of 5-litre containers of milk he will sell. (2)
- 2.4 Refer to the correlation between the price of 5-litre containers of milk and the number of 5-litre containers of milk sold, and comment on the accuracy of your answer to QUESTION 2.3. (2)
[10]



QUESTION 3

In the diagram, $A(-2 ; 10)$, $B(k ; k)$ and $C(4 ; -2)$ are the vertices of $\triangle ABC$. Line BC is produced to H and cuts the x -axis at $E(12 ; 0)$. AB and AC intersect the x -axis at F and G respectively. The angle of inclination of line AB is $81,87^\circ$.



3.1 Calculate the gradient of:

3.1.1 BE (2)

3.1.2 AB (2)

3.2 Determine the equation of BE in the form $y = mx + c$ (2)

3.3 Calculate the:

3.3.1 Coordinates of B , where $k < 0$ (2)



3.3.2 Size of \hat{A} (4)

3.3.3 Coordinates of the point of intersection of the diagonals of parallelogram $ACES$, where S is a point in the first quadrant (2)

3.4 Another point $T(p ; p)$, where $p > 0$, is plotted such that $ET = BE = 4\sqrt{17}$ units.

3.4.1 Calculate the coordinates of T . (5)

3.4.2 Determine the equation of the:


(a) Circle with centre at E and passing through B and T in the form $(x - a)^2 + (y - b)^2 = r^2$ (2)

(b) Tangent to the circle at point $B(k ; k)$ (3)

[24]

QUESTION 4

In the diagram, the circle centred at $N(-1 ; 3)$ passes through $A(-1 ; -1)$ and C . $B(-4 ; 2)$, C , D and E are joined to form a parallelogram such that BE is parallel to the x -axis. CD is a tangent to the circle at C and $CD = 6$ units.



- 4.1 Write down the length of the radius of the circle. (1)
- 4.2 Calculate the:
- 4.2.1 Coordinates of C (2)
 - 4.2.2 Coordinates of D (2)
 - 4.2.3 Area of $\Delta ABCD$ (3)
- 4.3 The circle, centred at N , is reflected about the line $y = x$. M is the centre of the new circle which is formed. The two circles intersect at A and F .
- Calculate the:
- 4.3.1 Length of NM (3)
 - 4.3.2 Midpoint of AF (4)

[15]

QUESTION 5

- 5.1 Without using a calculator, simplify the following expression to ONE trigonometric ratio:

$$\frac{\sin 140^\circ \cdot \sin(360^\circ - x)}{\cos 50^\circ \cdot \tan(-x)} \quad (6)$$

- 5.2 Prove the identity: $\frac{-2\sin^2 x + \cos x + 1}{1 - \cos(540^\circ - x)} = 2\cos x - 1 \quad (4)$

- 5.3 Given: $\sin 36^\circ = \sqrt{1 - p^2}$

Without using a calculator, determine EACH of the following in terms of p :

5.3.1 $\tan 36^\circ \quad (3)$

5.3.2 $\cos 108^\circ \quad (4)$

[17]

QUESTION 6

- 6.1 Given: $\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$

- 6.1.1 Use the given identity to derive a formula for $\cos(\alpha + \beta) \quad (3)$

- 6.1.2 Simplify completely: $2\cos 6x \cos 4x - \cos 10x + 2\sin^2 x \quad (5)$


- 6.2 Determine the general solution of $\tan x = 2 \sin 2x$ where $\cos x < 0. \quad (7)$

[15]



QUESTION 7

In the diagram below, the graph of $f(x) = -2 \sin x$ is drawn for the interval $x \in [-180^\circ; 180^\circ]$.




- 7.1 On the grid provided in the ANSWER BOOK, draw the graph of $g(x) = \cos(x - 60^\circ)$ for $x \in [-180^\circ; 180^\circ]$. Clearly show ALL intercepts with the axes and turning points of the graph. (3)
- 7.2 Write down the period of $f(3x)$. (2)
- 7.3 Use the graphs to determine the value of x in the interval $x \in [-180^\circ; 180^\circ]$ for which $f(x) - g(x) = 1$. (1)
- 7.4 Write down the range of k , if $k(x) = \frac{1}{2}g(x) + 1$. (2)
[8]



QUESTION 8

In the diagram below, T is a hook on the ceiling of an art gallery. Points Q, S and R are on the same horizontal plane from where three people are observing the hook T. The angle of elevation from Q to T is x .


$\hat{QSR} = 90^\circ + x$, $\hat{QRS} = x$, $QR = 5$ units and $TS = SQ$.



- 8.1 Prove that $QS = 5 \tan x$ (3)
- 8.2 Prove that the length of $QT = 10 \sin x$ (5)
- 8.3 Calculate the area of ΔTQR if $\hat{TQR} \approx 70^\circ$ and $x = 25^\circ$. (2)
[10]

QUESTION 9


In the diagram, PQRS is a cyclic quadrilateral. PS is produced to W. TR and TS are tangents to the circle at R and S respectively. $\hat{T} = 78^\circ$ and $\hat{Q} = 93^\circ$.



- 9.1 Give a reason why $ST = TR$. (1)
- 9.2 Calculate, giving reasons, the size of:
- 9.2.1 \hat{S}_2 (2)
- 9.2.2 \hat{S}_3 (2)
[5]

QUESTION 10


In the diagram, BE and CD are diameters of a circle having M as centre. Chord AE is drawn to cut CD at F. $AE \perp CD$. Let $\hat{C} = x$.



- 10.1 Give a reason why $AF = FE$. (1)
- 10.2 Determine, giving reasons, the size of \hat{M}_1 in terms of x . (3)
- 10.3 Prove, giving reasons, that AD is a tangent to the circle passing through A, C and F. (4)
- 10.4 Given that $CF = 6$ units and $AB = 24$ units, calculate, giving reasons, the length of AE. (5)
[13]


QUESTION 11

- 11.1 In the diagram, chords DE , EF and DF are drawn in the circle with centre O .
 KFC is a tangent to the circle at F .



Prove the theorem which states that $\hat{DFK} = \hat{E}$. (5)

- 11.2 In the diagram, PK is a tangent to the circle at K. Chord LS is produced to P. N and M are points on KP and SP respectively such that $MN \parallel SK$. Chord KS and LN intersect at T.



11.2.1 Prove, giving reasons, that:

(a) $\hat{K}_4 = \hat{NML}$ (4)

(b) KLMN is a cyclic quadrilateral (1)

11.2.2 Prove, giving reasons, that $\triangle LKN \parallel\!\!\!\parallel \triangle KSM$. (5)

11.2.3 If $LK = 12$ units and $3KN = 4SM$, determine the length of KS. (4)

11.2.4 If it is further given that $NL = 16$ units, $LS = 13$ units and $KN = 8$ units, determine, with reasons, the length of LT. (4)

[23]

TOTAL: 150

INFORMATION SHEET: MATHEMATICS

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$A = P(1 + ni)$$

$$A = P(1 - ni)$$

$$A = P(1 - i)^n$$

$$A = P(1 + i)^n$$

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2} [2a + (n - 1)d]$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(r^n - 1)}{r - 1}; \quad r \neq 1$$

$$S_\infty = \frac{a}{1 - r}; \quad -1 < r < 1$$

$$F = \frac{x[(1+i)^n - 1]}{i}$$

$$P = \frac{x[1 - (1+i)^{-n}]}{i}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$M\left(\frac{x_1 + x_2}{2}; \frac{y_1 + y_2}{2}\right)$$

$$y = mx + c$$

$$y - y_1 = m(x - x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \tan \theta$$

$$(x - a)^2 + (y - b)^2 = r^2$$

$$\text{In } \Delta ABC: \quad \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\text{area } \Delta ABC = \frac{1}{2} ab \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos 2\alpha = \begin{cases} \cos^2 \alpha - \sin^2 \alpha \\ 1 - 2\sin^2 \alpha \\ 2\cos^2 \alpha - 1 \end{cases}$$

$$\sin 2\alpha = 2\sin \alpha \cos \alpha$$

$$\bar{x} = \frac{\sum x}{n}$$

$$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

$$P(A) = \frac{n(A)}{n(S)}$$

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

$$\hat{y} = a + bx$$

$$b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$





NSC Answer Book
NSS-antwoordeboek

National Senior Certificate/Nasionale Senior Sertifikaat (Grade 12/Graad 12) – November 2021

CENTRE NUMBER SENTRUMNOMMER							
--------------------------------	--	--	--	--	--	--	--

EXAMINATION NUMBER EKSAMENNOMMER											
-------------------------------------	--	--	--	--	--	--	--	--	--	--	--

DATE DATUM												BOOK NUMBER BOEKNOMMER		OF VAN		BOOKS BOEKIE
---------------	--	--	--	--	--	--	--	--	--	--	--	---------------------------	--	-----------	--	-----------------

SUBJECT CODE TAKKODE												PAPER NUMBER VRAESTELNOMMER	2
-------------------------	--	--	--	--	--	--	--	--	--	--	--	--------------------------------	---

SUBJECT NAME VAKNAAM	MATHEMATICS/WISKUNDE
-------------------------	----------------------

MARKER/NASTIENER			MODERATOR'S INITIALS IN RELEVANT BLOCK MODERATOR SE VOORLETTERS IN RELEVANTE BLOKKIE									
Question Vraag	Marks Punte	Marker's Code & Initials Nastener se kode & Voorletters	Marks Punte	SM	Marks Punte	DCM AHN	Marks Punte	CM HN	Marks Punte	IM	Marks Punte	EM
1												
2												
3												
4												
5												
6												
7												
8												
9												
10												
11												
TOTAL TOTAAL												

CONTROLLED AND CERTIFIED CORRECT
(SURNAME AND INITIALS OF EA)
GEKONTROLEER EN AS KORREK
GESERTIFISEER (VAN EN VOORLETTERS
VAN EA)

READ INSTRUCTIONS ON THE NEXT PAGE
LEES INSTRUKSJIES OP VOLgende BlAAS!

This answer book consists of 25 pages./Hierdie antwoordeboek bestaan uit 25 bladsye.



PLEASE FOLLOW THESE INSTRUCTIONS CAREFULLY	VOLG ASSEBLIEF HIERDIE INSTRUKSIES NOUKEURIG
<ol style="list-style-type: none"> 1. Clearly write your examination number and centre number in the space provided and attach your barcode label in the space provided. 2. Remember that your own name (or the name of your school) may NOT appear anywhere on or in this answer book. 3. Answer ALL questions in the spaces provided. 4. NO pages may be torn from this answer book. 5. Read the instructions printed on your timetable carefully as well as any other instructions which may be given in each examination paper. 6. Candidates may NOT retain an answer book or remove it from the examination room. 7. Answers must be written in black/blue ink as distinctly as possible. Do NOT write in the margins. 8. Write the numbers of the questions you have answered on the front cover of the answer book where marks are to be recorded. 9. If you require additional space for your answers: <ol style="list-style-type: none"> 9.1 Use the additional space provided at the end of the answer book. 9.2 When answering a question in the additional space, indicate clearly the question number in the column on the LHS. 9.3 Rule off after each answer. 10. Draw a neat line through any work/rough work that must NOT be marked. 	<ol style="list-style-type: none"> 1. Skryf jou eksamennummer en sentrumnommer duidelik in die ruimtes soos verskaf en plak jou stafieskodeplakker in die ruimte soos verskaf. 2. Onthou dat jou eie naam (of die naam van jou skool) NIE op of in hierdie antwoordeboek mag voorkom NIE. 3. Beantwoord ALLE vrae in die ruimtes wat voorsien is. 4. GEEN bladsye mag uit hierdie antwoordeboek geskeur word NIE. 5. Lees die instruksies wat op jou eksamenrooster gedruk is sorgvuldig deur, asook enige ander instruksies wat op elke eksamenvraestel gegee word. 6. GEEN antwoordeboek mag deur die kandidaat behou of uit die eksamenlokaal verwyder word NIE. 7. Skryf die antwoorde so duidelik moontlik met swart/blou ink. Laat die kanlyne oop. 8. Skryf die nommers van die vrae wat jy beantwoord het op die voorblad van die antwoordeboek waar die punte aangebring word. 9. In geval jy bykomende ruimte benodig vir jou antwoorde: <ol style="list-style-type: none"> 9.1 Gebruik die bykomende ruimte wat aan die einde van die antwoordeboek verskaf word. 9.2 As 'n vraag in die bykomende ruimte beantwoord word, dui duidelik die vraagnommer in die kolom aan die LK aan. 9.3 Trek 'n lyn na elke antwoord. 10. Trek 'n netjiese lyn deur enige werk/rofwerk wat NIE nagesien moet word NIE.



QUESTION/VRAAG 1


1.1

10	11	13	14	14	15	16	18	18
19	19	20	21	35	35	37	40	41

	Solution/Oplossing	Marks/Punte
1.1.1	 Stanmorephysics.com	(2)
1.1.2		(1)
1.1.3		(2)
1.2		(2)
1.3.1		(2)
1.3.2		(1)
		[10]


QUESTION/VRAAG 2

Price of milk in rands per 5-litre container (x)/ <i>Prys van melk, in rand, per 5 liter-houer (x)</i>	26	32	36	28	40	33	29	34	27	30
Number of 5-litre containers of milk sold (y)/ <i>Aantal 5 liter-houers melk verkoop (y)</i>	48	30	26	44	23	32	39	29	42	33




	Solution/Oplossing	Marks/Punte
2.2		
2.3		
2.4		
		[10]

QUESTION/VRAAG 3




	Solution/Oplossing	Marks/Punte
3.1.1		(2)
3.1.2		(2)
3.2		(2)
3.3.1		(2)

	Solution/ <i>Oplossing</i>	Marks/ <i>Punte</i>
3.3.2		(4)
3.3.3		(2)
3.4.1		(5)
3.4.2(a)		(2)
3.4.2(b)		(3)



QUESTION/VRAAG 4



	Solution/ <i>Oplossing</i>	Marks/ <i>Punte</i>
4.1		(1)
4.2.1		(2)
4.2.2		(2)
4.2.3		(3)

	Solution/Oplossing	Marks/Punte
4.3.1		(3)
4.3.2		(4) [15]

QUESTION/VRAAG 5

	Solution/Oplossing	Marks/Punte
5.1		(6)
5.2		(4)
5.3.1		(3)




5.3.2		
		(4) [17]

QUESTION/VRAAG 6

	Solution/ <i>Oplossing</i>	Marks/ <i>Punte</i>
6.1.1		(3)
6.1.2		(5)

	Solution/Oplossing	Marks/Punte
6.2		(7) [15]




QUESTION/VRAAG 7

	Solution/Oplossing	Marks/Punte
7.1		(3)
7.2		(2)
7.3		(1)
7.4		(2) [8]

Give reasons for your statements in QUESTIONS 8, 9, 10 and 11.
Gee redes vir jou bewerings in VRAAG 8, 9, 10 en 11.

QUESTION/VRAAG 8




	Solution/Oplossing	Marks/Punte
8.1	(3)
8.2	(5)

8.3		(2)	[10]
-----	--	-----	------




QUESTION/VRAAG 9



	Solution/Oplossing	Marks/Punte
9.1		(1)
9.2.1		(2)
9.2.2		(2) [5]


QUESTION/VRAAG 10



	Solution/Oplossing	Marks/Punte
10.1		(1)
10.2		(3)
10.3		(4)
10.4		(5)
		[13]


QUESTION/VRAAG 11

11.1



	Solution/Oplossing	Marks/Punte
		(5)

11.2




	Solution/Opllossing	Marks/Punte
11.2.1(a)		(4)
11.2.1(b)		(1)

	Solution/Oplossing	Marks/Punte
11.2.2		(5)
11.2.3		(4)
11.2.4		(4)

[23]



	Additional space/Bykomende ruimte	Marks/Punte



	Additional space/ <i>Bykomende ruimte</i>	Marks/ <i>Punte</i>




	Additional space/Bykomende ruimte	Marks/Punte



	Additional space/ <i>Bykomende ruimte</i>	Marks/ <i>Punte</i>

TOTAL/TOTAAL: 150



RE-MARK/RE-CHECK HERMERK/HERSIEN			
Question Vraag	Marks Punte	Initials Voorletters	
1			
2			
3			
4			
5			
6			
7			
8			
9			
10			
11			
TOTAL TOTAAL			
HASH TOTAL KAF- TOTAAL			





basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

NATIONAL SENIOR CERTIFICATE/ NASIONALE SENIOR SERTIFIKAAT

GRADE/GRAAD 12

MATHEMATICS P2/WISKUNDE V2

stanmorephysics.com

NOVEMBER 2021

MARKING GUIDELINES/NASIENRIGLYNE

MARKS/PUNTE: 150

These marking guidelines consist of 24 pages.
Hierdie nasienriglyne bestaan uit 24 bladsye.

NOTE:

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed out version.
- Consistent accuracy applies in ALL aspects of the marking memorandum. Stop marking at the second calculation error.
- Assuming answers/values in order to solve a problem is NOT acceptable.


NOTA:

- As 'n kandidaat 'n vraag TWEE KEER beantwoord, sien slegs die EERSTE poging na.
- As 'n kandidaat 'n antwoord van 'n vraag doodtrek en nie oordoen nie, sien die doodgetrekte poging na.
- Volgehoue akkuraatheid word in ALLE aspekte van die nasienriglyne toegepas. Hou op nasien by die tweede berekeningsfout.
- Om antwoorde/waardes te aanvaar om 'n probleem op te los, word NIE toegelaat NIE.

GEOMETRY • MEETKUNDE	
S	A mark for a correct statement (A statement mark is independent of a reason)
	<i>'n Punt vir 'n korrekte bewering</i> <i>('n Punt vir 'n bewering is onafhanklik van die rede)</i>
R	A mark for the correct reason (A reason mark may only be awarded if the statement is correct)
	<i>'n Punt vir 'n korrekte rede</i> <i>('n Punt word slegs vir die rede toegeken as die bewering korrek is)</i>
S/R	Award a mark if statement AND reason are both correct
	<i>Ken 'n punt toe as die bewering EN rede beide korrek is</i>

QUESTION/VRAAG 1

10	11	13	14	14	15	16	18	18
19	19	20	21	35	35	37	40	41




1.1.1	$\bar{x} = \frac{396}{18}$ $\bar{x} = 22$	Answer only: Full marks <i>Slegs antw: Volpunte</i>	\checkmark 396 \checkmark answer (2)
1.1.2	$\sigma = 10,1707 \approx 10,17$		\checkmark answer (1)
1.1.3	$\bar{x} + \sigma = 32,17$ $\therefore 5$ days	\checkmark 32,17 \checkmark 5	(2)
1.2	$22 \times 18 = 396$ ordered/bestel $20 \times 18 = 360$ sold/verkoop Total not sold/Totaal nie verkoop nie: 36 OR/OF $22 - 20 = 2$ $2 \times 18 = 36$	\checkmark $18\bar{x}_1$ and $18\bar{x}_2$ \checkmark answer	(2)
1.3.1	Option B/Opsie B <u>Any one of the following reasons/Enige een van die vlg redes:</u> <ul style="list-style-type: none"> • Median/Mediaan = 18,5 • $Q_1 = 14$ • IQR = 21 • Mean > Median, therefore the data is skewed to the right 	\checkmark B \checkmark reason	(2)
1.3.2	Data is positively skewed/skewed to the right <i>Data is positief skeef/skeef na regs</i>	\checkmark answer	(1)
[10]			

QUESTION/VRAAG 2

Price of milk in rands per 5-litre container (x) <i>Prys van melk in rand, per 5 liter-houer (x)</i>	26	32	36	28	40	33	29	34	27	30
Number of 5-litre containers of milk sold (y) <i>Aantal 5 liter-houers melk verkoop (y)</i>	48	30	26	44	23	32	39	29	42	33

2.1	<p style="text-align: center;">SCATTER PLOT</p> <table border="1"> <thead> <tr> <th>Price per 5 litre</th> <th>Units sold</th> </tr> </thead> <tbody> <tr><td>26</td><td>48</td></tr> <tr><td>27</td><td>42</td></tr> <tr><td>28</td><td>39</td></tr> <tr><td>29</td><td>33</td></tr> <tr><td>30</td><td>30</td></tr> <tr><td>31</td><td>29</td></tr> <tr><td>32</td><td>27</td></tr> <tr><td>33</td><td>26</td></tr> <tr><td>34</td><td>24</td></tr> </tbody> </table>	Price per 5 litre	Units sold	26	48	27	42	28	39	29	33	30	30	31	29	32	27	33	26	34	24	1 mark: 3 to 5 points plotted correctly 2 marks: 6 to 9 points plotted correctly 3 marks: all points plotted correctly (3)
Price per 5 litre	Units sold																					
26	48																					
27	42																					
28	39																					
29	33																					
30	30																					
31	29																					
32	27																					
33	26																					
34	24																					
2.2	$a = 90,478\dots \approx 90,48$ $b = -1,773\dots \approx -1,77$ $\hat{y} = 90,48 - 1,77x$	<div style="border: 1px solid black; padding: 5px;"> Answer only: Full marks <i>Slegs antw: Volpunte</i> </div>	✓ a ✓ b ✓ equation (3)																			
2.3	$y = 23,069\dots \approx 23,07$ units/eenhede (calculator/sakrekenaar) OR/OF $y = 90,48 - 1,77(38)$ $y = 23,22$ units/eenhede	✓✓ answer ✓ substitution ✓ answer (2)																				
2.4	$r = -0,94$ The value of r indicates a strong relationship between the cost per 5 litre and the number of units sold \therefore there is a good chance of the prediction being accurate. <i>Die waarde van r dui 'n sterke verwantskap tussen die koste per 5 liter en die aantal eenhede verkoop aan \therefore daar is 'n goeie kans dat die voorspelling akkuraat is</i>	✓ value of r OR/OF strong relationship/ <i>sterk verwantskap</i> ✓ accurate/akkuraat (2)																				
[10]																						


QUESTION/VRAAG 3



<p>3.3.1</p> $y = \frac{1}{4}x - 3$ $k = \frac{1}{4}k - 3$ $\frac{3}{4}k = -3$ $k = -4$ $\therefore B(-4; -4)$ <p>OR/OF</p> $m_{BE} = \frac{1}{4}$ $\frac{0-k}{12-k} = \frac{1}{4}$ $-4k = 12 - k$ $k = -4$ $\therefore B(-4; -4)$ <p>OR/OF</p> $m_{AB} = \tan 81,87^\circ$ $m_{AB} = 7$ $m_{AB} = \frac{10-k}{-2-k}$ $7(-2-k) = 10 - k$ $-14 - 7k = 10 - k$ $-6k = 24$ $k = -4$ $\therefore B(-4; -4)$ <p>OR/OF</p> $\text{EB: } y = \frac{1}{4}x - 3 \quad \text{and AB: } y = 7x + 24$ $\frac{1}{4}x - 3 = 7x + 24$ $\frac{27}{4}x = -27$ $x = k = -4$ $\therefore B(-4; -4)$	<p>✓ substitution</p> <p>✓ answer (2)</p> <p>✓ substitution</p> <p>✓ answer (2)</p> <p>✓ substitution</p> <p>✓ answer (2)</p> <p>✓ equating EB & AB</p> <p>✓ answer (2)</p>
---	---

3.3.2	<p>In ΔAFG:</p> $m_{AC} = \frac{10 - (-2)}{-2 - 4} = -2$ $\tan \theta = m_{AC} = -2$ $\theta = 180^\circ - 63,43\dots^\circ$ $\therefore \theta = 116,57^\circ$ $\therefore \hat{A} = 116,57^\circ - 81,87^\circ \text{ [ext } \angle \text{ of } \Delta \text{]}$ $\therefore \hat{A} = 34,70^\circ$ <p>OR/OF</p> <p>In ΔABC:</p> $a = BC = 2\sqrt{17}; b = AC = 6\sqrt{5}; c = AB = 10\sqrt{2}$ $a^2 = b^2 + c^2 - 2bc \cdot \cos A$ $(2\sqrt{17})^2 = (6\sqrt{5})^2 + (10\sqrt{2})^2 - 2(6\sqrt{5})(10\sqrt{2}) \cdot \cos A$ $\cos A = \frac{(6\sqrt{5})^2 + (10\sqrt{2})^2 - (2\sqrt{17})^2}{2(6\sqrt{5})(10\sqrt{2})}$ $= 0,822\dots$ $\therefore A = 34,7^\circ$	$\checkmark m_{AC} = -2$ $\checkmark \tan \theta = -2$ $\checkmark \theta = 116,57^\circ$ \checkmark answer (4)
3.3.3	$M\left(\frac{12 + (-2)}{2}; \frac{10 + (0)}{2}\right)$ Diagonals intersect at the point (5 ; 5)	\checkmark x-value \checkmark y-value (2)
3.4.1	$BE = ET$ $4\sqrt{17} = \sqrt{(12 - p)^2 + (0 - p)^2}$ $(4\sqrt{17})^2 = (\sqrt{(12 - p)^2 + (0 - p)^2})^2$ $272 = 144 - 24p + p^2 + p^2$ $p^2 - 12p - 64 = 0$ $(p - 16)(p + 4) = 0$ $\therefore p = 16 \quad \text{or} \quad p = -4 \text{ (n.a.)}$ $\therefore T(16; 16)$	\checkmark substitution of E & T \checkmark equating \checkmark standard form \checkmark factors $\checkmark p = 16$ (5)
3.4.2a	$(x - 12)^2 + y^2 = (4\sqrt{17})^2 = 272$	\checkmark LHS \checkmark RHS (2)
3.4.2b	$m_{radius} = \frac{1}{4}$ $m_{tangent} = -4$ $y = -4x + c$ OR/OF $y - y_1 = -4(x - x_1)$ $-4 = -4(-4) + c$ $y - (-4) = -4(x - (-4))$ $c = -20$ $y = -4x - 20$ $y = -4x - 20$	$\checkmark m_{tangent}$ \checkmark substitution of B \checkmark equation (3) [24]



QUESTION/VRAAG 4



4.1	Radius = 4 units/eenhede	✓ answer (1)
4.2.1	CD \perp CN $\therefore C(-1; 7)$	✓ x value ✓ y value (2)
4.2.2	CD = 6 units $\therefore D(5; 7)$	✓ x value ✓ y value (2)
4.2.3	$\perp h = 5$ units DC = 6 units Area $\Delta ABCD = \frac{1}{2}(6)(5)$ $= 15$ units ²	✓ $\perp h = 5$ units ✓ substitution into Area formula ✓ answer (3)
OR/OF		
	$\perp h = 5$ units DC = 6 units Area $\Delta ABCD = \frac{1}{2}[\text{Area of } ^m]$ $= \frac{1}{2}[(5)(6)]$ $= 15$ units ²	✓ $\perp h = 5$ units ✓ substitution into Area formula ✓ answer (3)

	<p>OR/OF</p> <p>Let angle of inclination of BC = α</p> $\tan \alpha = \frac{5}{3}$ $\alpha = 59,036\dots^\circ$ $\hat{B}CD = 180^\circ - \alpha$ $\hat{B}CD = 180^\circ - 59,036\dots^\circ$ $\hat{B}CD = 120,96^\circ$ $\text{Area } \Delta ABCD = \frac{1}{2}(\sqrt{34})(6)\sin 120,96^\circ$ $= 15 \text{ units}^2$	
4.3.1	<p>M(3 ; -1) [reflection of N(-1 ; 3) about the line $y = x$]</p> $\therefore MN = \sqrt{(3 - (-1))^2 + (-1 - 3)^2}$ $MN = \sqrt{32} = 4\sqrt{2} = 5,66 \text{ units}$	✓ coordinates of M (A) ✓ substitution of M&N ✓ answer (3)
4.3.2	<p>M(3 ; -1)</p> $m_{MN} = \frac{3 - (-1)}{-1 - 3} = -1$ <p>MN: $-1 = -(3) + c$ or $y - 3 = -1(x + 1)$ $c = 2$ $y - 3 = -x - 1$ $\therefore y = -x + 2$ $y = -x + 2$</p> <p>$x = -x + 2$ $2x = 2$ $x = 1$ $\therefore y = 1$ midpoint (1 ; 1)</p>	✓ equation of MN ✓ equating AF & MN ✓ x value ✓ y value (4)
	<p>OR/OF</p> <p>N(-1 ; 3)</p> $y_F = y_N = 3$ <p>Reflected about $y = x$</p> $\therefore F(3 ; 3)$ $\text{midpoint}\left(\frac{-1+3}{2}; \frac{-1+3}{2}\right) = (1 ; 1)$	<p>✓✓ coordinates of F ✓ x value ✓ y value (4)</p>

OR/OF



NAMF is a square ($NA=NF=AM=MF$ and $NA \perp AM$)

$$\begin{aligned} \text{Midpoint NM} &= (1 ; 1) \\ &= \text{Midpoint of AF} \end{aligned}$$


✓ NAMF = square

✓ x ✓ y of midpt NM
 ✓ midpt AF

(4)

[15]

QUESTION/VRAAG 5

<p>5.1</p> $ \begin{aligned} & \frac{\sin 140^\circ \cdot \sin(360^\circ - x)}{\cos 50^\circ \cdot \tan(-x)} \\ &= \frac{\sin 40^\circ (-\sin x)}{\sin 40^\circ (-\tan x)} \\ &= \frac{-\sin x}{-\frac{\sin x}{\cos x}} \\ &= \cos x \end{aligned} $	<ul style="list-style-type: none"> ✓ $\sin 40^\circ$ ✓ $-\sin x$ ✓ co-ratio ✓ $-\tan x$ ✓ $\tan x = \frac{\sin x}{\cos x}$ ✓ answer
<p>5.2</p> $ \begin{aligned} \text{LHS} &= \frac{-2\sin^2 x + \cos x + 1}{1 - \cos(540^\circ - x)} & \text{RHS} &= 2\cos x - 1 \\ \text{LHS} &= \frac{-2(1 - \cos^2 x) + \cos x + 1}{1 - (-\cos x)} \\ \text{LHS} &= \frac{-2 + 2\cos^2 x + \cos x + 1}{1 + \cos x} \\ \text{LHS} &= \frac{2\cos^2 x + \cos x - 1}{1 + \cos x} \\ \text{LHS} &= \frac{(2\cos x - 1)(\cos x + 1)}{1 + \cos x} \\ \text{LHS} &= 2\cos x - 1 \\ \therefore \text{LHS} &= \text{RHS} \end{aligned} $	<ul style="list-style-type: none"> ✓ identity i. t. o. $\cos x$ ✓ $\cos(540^\circ - x) = -\cos x$ ✓ standard form ✓ factors
<p>5.3.1</p> $ \begin{aligned} \sin 36^\circ &= \sqrt{1 - p^2} \\ \tan 36^\circ &= \frac{\sqrt{1 - p^2}}{p} \end{aligned} $ <p>OR/OF</p> $ \begin{aligned} \cos^2 36^\circ &= 1 - \sin^2 36^\circ \\ \cos 36^\circ &= \sqrt{1 - (1 - p^2)} \\ &= p \end{aligned} $ $ \begin{aligned} \tan 36^\circ &= \frac{\sin 36^\circ}{\cos 36^\circ} \\ &= \frac{\sqrt{1 - p^2}}{p} \end{aligned} $	 <ul style="list-style-type: none"> ✓ method ✓ value of p ✓ answer

<p>5.3.2</p> $\begin{aligned} &\cos 108^\circ \\ &= -\cos 72^\circ \\ &= -\cos(2 \times 36^\circ) \\ &= -(2 \cos^2 36^\circ - 1) \\ &= -2p^2 + 1 \end{aligned}$ <p>OR/OF</p> $\begin{aligned} &\cos 108^\circ \\ &= -\cos 72^\circ \\ &= -\cos(2 \times 36^\circ) \\ &= -(1 - 2 \sin^2 36^\circ) \\ &= -1 + 2(\sqrt{1 - p^2})^2 \\ &= -1 + 2(1 - p^2) \\ &= -2p^2 + 1 \end{aligned}$ <p>OR/OF</p> $\begin{aligned} &\cos 108^\circ \\ &= -\cos 72^\circ \\ &= -\cos(2 \times 36^\circ) \\ &= -(\cos^2 36^\circ - \sin^2 36^\circ) \\ &= -(p^2 - (\sqrt{1 - p^2})^2) \\ &= -(p^2 - (1 - p^2)) \\ &= -2p^2 + 1 \end{aligned}$ <p>OR/OF</p> $\begin{aligned} &\cos 108^\circ \\ &= \cos(2 \times 54^\circ) \\ &= 2 \cos^2 54^\circ - 1 \\ &= 2(1 - p^2) - 1 \\ &= 1 - 2p^2 \end{aligned}$ <p>OR/OF</p> $\begin{aligned} \cos 108^\circ &= \cos(72^\circ + 36^\circ) \\ &= \cos 72^\circ \cos 36^\circ - \sin 72^\circ \sin 36^\circ \\ &= (2 \cos^2 36^\circ - 1) \cos 36^\circ - (2 \sin 36^\circ \cos 36^\circ) \sin 36^\circ \\ &= 2 \cos^3 36^\circ - \cos 36^\circ - 2 \cos 36^\circ \sin^2 36^\circ \\ &= 2p^3 - p - 2p(\sqrt{1 - p^2})^2 \\ &= 2p^3 - p - 2p + 2p^3 \\ &= 4p^3 - 3p \end{aligned}$	<ul style="list-style-type: none"> ✓ reduction ✓ double angle ✓ expansion ✓ answer i. t. o. p (4) <ul style="list-style-type: none"> ✓ reduction ✓ double angle ✓ expansion ✓ answer i. t. o. p (4) <ul style="list-style-type: none"> ✓ reduction ✓ double angle ✓ expansion ✓ answer i. t. o. p (4) <ul style="list-style-type: none"> ✓ double angle ✓ expansion ✓ answer i. t. o. p (4) <ul style="list-style-type: none"> ✓ expansion ✓ both double angle identities ✓ value of $\sin 36^\circ$ ✓ answer i. t. o. p (4)
	[17]


QUESTION/VRAAG 6

6.1.1	$\begin{aligned} & \cos(\alpha + \beta) \\ &= \cos(\alpha - (-\beta)) \\ &= \cos \alpha \cos(-\beta) + \sin \alpha \sin(-\beta) \\ &= \cos \alpha \cos \beta + \sin \alpha (-\sin \beta) \\ &= \cos \alpha \cos \beta - \sin \alpha \sin \beta \end{aligned}$	<ul style="list-style-type: none"> ✓ $\cos(\alpha - (-\beta))$ ✓ expansion ✓ reduction 	(3)
6.1.2	$\begin{aligned} & 2 \cos 6x \cos 4x - \cos 10x + 2 \sin^2 x \\ &= 2 \cos 6x \cos 4x - \cos(6x + 4x) + 2 \sin^2 x \\ &= 2 \cos 6x \cos 4x - (\cos 6x \cos 4x - \sin 6x \sin 4x) + 2 \sin^2 x \\ &= \cos 6x \cos 4x + \sin 6x \sin 4x + 2 \sin^2 x \\ &= \cos 2x + 2 \sin^2 x \\ &= 1 - 2 \sin^2 x + 2 \sin^2 x \\ &= 1 \end{aligned}$	<ul style="list-style-type: none"> ✓ $\cos 10x = \cos(6x + 4x)$ ✓ expansion of $\cos(6x + 4x)$ ✓ $\cos 2x$ ✓ $1 - 2 \sin^2 x$ ✓ answer 	(5)
6.2	$\begin{aligned} \tan x &= 2 \sin 2x \\ \frac{\sin x}{\cos x} &= 2(2 \sin x \cos x) \\ \sin x &= 4 \sin x \cos^2 x \\ 4 \sin x \cos^2 x - \sin x &= 0 \\ \sin x(4 \cos^2 x - 1) &= 0 \\ \sin x = 0 & \quad \text{or} \quad \cos^2 x = \frac{1}{4} \\ & \quad \cos x = -\frac{1}{2} \\ x = 180^\circ + k \cdot 360^\circ; k \in \mathbb{Z} & \quad \text{or} \quad x = 120^\circ + k \cdot 360^\circ; k \in \mathbb{Z} \\ & \quad x = 240^\circ + k \cdot 360^\circ; k \in \mathbb{Z} \end{aligned}$ <p>OR/OF</p> $\begin{aligned} \tan x &= 2 \sin 2x \\ \frac{\sin x}{\cos x} &= 4 \sin x \cos x \\ \sin x &= 4 \sin x \cos^2 x \\ 4 \sin x \cos^2 x - \sin x &= 0 \\ 4 \sin x(1 - \sin^2 x) - \sin x &= 0 \\ 3 \sin x - 4 \sin^3 x &= 0 \\ \sin x(3 - 4 \sin^2 x) &= 0 \\ \sin x = 0 & \quad \text{or} \quad \sin^2 x = \frac{3}{4} \\ & \quad \sin x = \frac{\sqrt{3}}{2} \quad \text{or} \quad \sin x = -\frac{\sqrt{3}}{2} \\ x = 180^\circ + k \cdot 360^\circ, k \in \mathbb{Z} & \quad \text{or} \quad x = 120^\circ + k \cdot 360^\circ, k \in \mathbb{Z} \\ & \quad \text{or} \quad x = 240^\circ + k \cdot 360^\circ, k \in \mathbb{Z} \end{aligned}$	<ul style="list-style-type: none"> ✓ quotient identity ✓ double angle identity ✓ factors ✓ both equations ✓ $x = 180^\circ$ ✓ $x = 120^\circ \& 240^\circ$ OR/OF $x = \pm 120^\circ$ ✓ $k \cdot 360^\circ; k \in \mathbb{Z}$ ✓ quotient identity ✓ identity ✓ factors ✓ both equations ✓ $x = 180^\circ$ ✓ $x = 120^\circ \& 240^\circ$ OR/OF $x = \pm 120^\circ$ ✓ $k \cdot 360^\circ; k \in \mathbb{Z}$ 	(7)
			[15]

QUESTION/VRAAG 7

7.1		<ul style="list-style-type: none"> ✓ both turning points ✓ both x intercepts (-30° & 150°) ✓ shape
7.2	Period = 120°	✓✓ answer (2)
7.3	$x = -30^\circ$	✓ answer (1)
7.4	<p>Range of/waardeversameling van g: $y \in [-1; 1]$</p> <p>Range of/Waardeversameling van $\frac{1}{2}g$: $y \in \left[-\frac{1}{2}; \frac{1}{2}\right]$</p> <p>Range of/Waardeversameling van $\frac{1}{2}g + 1$: $y \in \left[\frac{1}{2}; \frac{3}{2}\right]$</p> <p>OR/OF</p> <p>Range of/Waardeversameling van $\frac{1}{2}g + 1$: $\frac{1}{2} \leq y \leq \frac{3}{2}$</p>	<ul style="list-style-type: none"> ✓ critical values ✓ correct notation (2) <ul style="list-style-type: none"> ✓ critical values ✓ correct notation (2)
[8]		


QUESTION/VRAAG 8



8.1	<p>In ΔSQR:</p> $\frac{QS}{\sin x} = \frac{QR}{\sin(90^\circ + x)}$ $\frac{QS}{\sin x} = \frac{5}{\cos x}$ $QS = \frac{5 \sin x}{\cos x}$ $QS = 5 \tan x$	<ul style="list-style-type: none"> ✓ correct use of sine rule ✓ $\sin(90^\circ + x) = \cos x$ ✓ $QS = \frac{5 \sin x}{\cos x}$ <p>(3)</p>
8.2	$\frac{QT}{\sin(180^\circ - 2x)} = \frac{TS}{\sin x}$ $\frac{QT}{\sin 2x} = \frac{5 \tan x}{\sin x}$ $QT = \frac{5 \tan x \sin 2x}{\sin x}$ $QT = \frac{5 \left(\frac{\sin x}{\cos x} \right) (2 \sin x \cos x)}{\sin x}$ $QT = \frac{5 \sin x (2 \sin x)}{\sin x}$ $QT = 10 \sin x$	<ul style="list-style-type: none"> ✓ correct use of sine rule ✓ $TS = QS = 5 \tan x$ ✓ $QT = \frac{5 \tan x \sin 2x}{\sin x}$ ✓ $\tan x = \frac{\sin x}{\cos x}$ ✓ $\sin 2x = 2 \sin x \cos x$ <p>(5)</p>

	<p>OR/OF</p> $QT^2 = QS^2 + TS^2 - 2QS \cdot TS \cos x$ $(5 \tan x)^2 = (5 \tan x)^2 + TQ^2 - 2(5 \tan x) \cdot TQ \cos x$ $0 = TQ^2 - 2(5 \tan x) \cdot TQ \cos x$ $0 = TQ [TQ - 10 \tan x \cos x]$ $TQ = 10 \tan x \cos x \quad (TQ \neq 0)$ $= 10 \frac{\sin x}{\cos x} \cdot \cos x$ $= 10 \sin x$	<ul style="list-style-type: none"> ✓ correct use of cos rule ✓ $TS = QS = 5 \tan x$ ✓ quadratic equation into TQ ✓ $TQ = 10 \tan x \cdot \cos x$ ✓ $\tan x = \frac{\sin x}{\cos x}$ <p>(5)</p>
8.3	<p>Area of $\Delta TQR = \frac{1}{2} \cdot TQ \cdot QR \sin T\hat{Q}R$</p> $= \frac{1}{2} (10 \sin 25^\circ)(5)(\sin 70^\circ)$ $= 9,93 \text{ unit}^2$	<ul style="list-style-type: none"> ✓ correct substitution into the area rule ✓ answer <p>(2)</p>
[10]		


QUESTION/VRAAG 9



9.1	tangents from same(common) point/raaklyne vanaf dieselfde punt	✓ R (1)
9.2.1	$\hat{S}_2 = \hat{S}RT$ $\therefore \hat{S}_2 = 51^\circ$	[\angle s opp equal sides/ \angle e teenoor gelyke sye] [sum of \angle s in Δ /som van \angle e in Δ] ✓ R ✓ S (2)
9.2.2	$\hat{S}_2 + \hat{S}_3 = 93^\circ$ $\hat{S}_3 = 42^\circ$ OR/OF $\hat{S}_1 = 87^\circ$ $\hat{S}_3 = 180^\circ - (87^\circ + 51^\circ)$ $\hat{S}_3 = 42^\circ$	[ext \angle of cyclic quad/buite \angle van koordevh] ✓ R ✓ answer ✓ R ✓ answer (2)

[5]

QUESTION/VRAAG 10




10.1	line from centre \perp to chord/lyn vanaf middelpunt \perp op koord	\checkmark R (1)
10.2	$\therefore \hat{A}_1 = 90^\circ - x$ [sum of \angle s in Δ /som van \angle e in Δ] $\therefore \hat{M}_1 = 180^\circ - 2x$ [\angle at centre = $2 \times$ \angle at circumf/midpts \angle = $2 \times$ omtreks \angle]	\checkmark S \checkmark S \checkmark R (3)
10.3	$\hat{CAD} = 90^\circ$ [\angle in semi circle/ \angle in halfsirkel] $\hat{A}_2 = 90^\circ - (90^\circ - x)$ $\hat{A}_2 = x$ $\therefore \hat{A}_2 = \hat{C} = x$ $\therefore AD$ is a tangent [converse tan-chord theorem/omgek rkl-kd st.]	\checkmark S \checkmark R \checkmark S \checkmark R (4)
OR/OF		
	$\hat{EMD} = 2x$ [adj suppl \angle s/aanligg suppl \angle e] $\therefore \hat{A}_2 = x$ [\angle at centre = $2 \times$ \angle at circumf/midpts \angle = $2 \times$ omtreks \angle] $\therefore \hat{A}_2 = \hat{C} = x$ $\therefore AD$ is a tangent [converse tan-chord theorem/omgek rkl-kd st.]	\checkmark S \checkmark S \checkmark R \checkmark R (4)
OR/OF		
	$\hat{M}_3 = 180^\circ - 2x$ [vert. opp/regoorstaande \angle e] $\therefore \hat{A}_3 = 90^\circ - x$ [\angle at centre = $2 \times$ \angle at circumf/midpts \angle = $2 \times$ omtreks \angle] $\hat{BAE} = 90^\circ$ [\angle in semi-circle/ \angle in halfsirkel] $\therefore \hat{A}_2 = \hat{C} = x$ $\therefore AD$ is a tangent [converse tan-chord theorem/omgek rkl-kd st.]	\checkmark S \checkmark R \checkmark S \checkmark R (4)

	<p>CD AB [midpt. Thm/ middelpuntst.] $\hat{B}AE = 90^\circ$ [\angle in semi-circle/\angle in halfsirkel] $\therefore \hat{A}_3 = \hat{D} = 90^\circ - x$ [alt.\angles; CD AB/verwiss \anglee] $\therefore \hat{A}_2 = x = C$ $\therefore AD$ is a tangent [converse tan-chord theorem/omgek rkl-kd st.]</p> <p>OR/OF</p> <p>$\hat{C}AD = 90^\circ$ [\angle in semi circle/\angle in halfsirkel] AC = diameter [converse \angle in semi circle/omgek \angle in halfsirkel] $\therefore AD$ is a tangent [converse radius \perp tangent/omgek radius \perp rkl]</p>	<ul style="list-style-type: none"> ✓ S ✓ R ✓ S ✓ R
10.4	<p>AF = FE and BM = ME [given & radii] $\therefore FM = \frac{1}{2} AB = 12$ units [Midpt Theorem/middelpuntstelling] $EM = MB = CM = 18$ units [radii] $\therefore EB = 36$ units [diameter = 2 radius] $\therefore AE^2 = (36)^2 - (24)^2$ [Pythagoras] $AE = 12\sqrt{5}$ or 26,83 units</p> <p>OR/OF</p> <p>AF = FE and BM = ME [given & radii] $\therefore FM = \frac{1}{2} AB = 12$ units [Midpt Theorem/middelpuntstelling] $EM = MB = CM = 18$ units [radii] $\therefore FE^2 = (18)^2 - (12)^2$ [Pythagoras] $FE = 6\sqrt{5}$ $AE = 12\sqrt{5}$ or 26,83 units</p>	<ul style="list-style-type: none"> ✓ FM = 12 ✓ R ✓ EB = 36 ✓ using Pyth correctly ✓ answer
		(4)
		(5)
		[13]


QUESTION/VRAAG 11

11.1




	<p>Construction: Draw diameter BF and draw BE <i>Konstruksie: Trek middellyn BF en verbind BE</i></p> $\hat{B}FK = 90^\circ \text{ or } \hat{DFK} = 90^\circ - \hat{BFD} \quad [\text{radius } \perp \text{tangent/raaklyn}]$ $\hat{BEF} = 90^\circ \quad [\angle \text{ in semi-circle/semi-sirkel}]$ $\therefore \hat{DEF} = 90^\circ - \hat{BED}$ $= 90^\circ - \hat{BFD} \quad [\angle \text{s same segment/}\angle \text{e dieselfde segment}]$ $\therefore \hat{DFK} = \hat{DEF}$	<p>✓ Constr ✓ S ✓ R ✓ S ✓ S/R</p>
		(5)

OR/OF




	<p>Construction: Draw radii DO and OF</p> <p><i>Konstruksie: Trek radii DO en OF</i></p> <p>$\hat{O}FK = 90^\circ$ or $\hat{D}FK = 90^\circ - \hat{O}FD$ [radius \perp tangent/<i>raaklyn</i>]</p> <p>$\hat{O}DF = \hat{O}FD$ [\angles opp = sides/\anglee teenoor = sye]</p> <p>$\therefore \hat{D}OF = 180^\circ - 2\hat{O}FD$ [\angles of Δ/\anglee van Δ]</p> <p>$\hat{D}EF = 90^\circ - \hat{O}FD$ [\angle at centre = $2 \times \angle$ circumf/ <i>midpts</i> \angle = $2 \times$ <i>omtreks</i> \angle]</p> <p>$\therefore \hat{D}FK = \hat{D}EF$</p>	<p>✓ construction</p> <p>✓ S ✓R</p> <p>✓ S</p> <p>✓ S/R</p> <p>(5)</p>
--	---	--

OR/OF



	<p>Construction: Draw diameter BF and join BD.</p> <p><i>Konstruksie: Trek middellyn BF en verbind BD.</i></p> <p>$\hat{B}FK = 90^\circ$ or $\hat{DFK} = 90^\circ - \hat{BFD}$ [radius \perp tangent/<i>raaklyn</i>]</p> <p>$\hat{FDB} = 90^\circ$ [\angle in half circle/<i>semi-sirkel</i>]</p> <p>$\hat{B} = 90^\circ - \hat{BFD}$</p> <p>$\therefore \hat{DFK} = \hat{B}$</p> <p>but $\hat{B} = \hat{E}$ [\angles same segment/<i>∠e dieselfde segment</i>]</p> <p>$\therefore \hat{DFK} = \hat{E}$</p>	<p>✓ construction</p> <p>✓ S ✓/R</p> <p>✓ S</p> <p>✓ S/R</p>	(5)
--	---	--	-----

11.2



11.2.1(a)	$\hat{K}_4 = \hat{S}_1$ [tan chord theorem / raaklynkoordstelling]	✓ S ✓ R
	$\hat{M}_2 + \hat{M}_3 = \hat{S}_1$ [corresp \angle s; / ooreenk \angle s; MN KS]	✓ S ✓ R
	$\therefore \hat{K}_4 = \hat{M}_2 + \hat{M}_3 = \hat{NML}$	(4)
11.2.1(b)	$\therefore \hat{K}_4 = \hat{M}_2 + \hat{M}_3 = \hat{NML}$	
	$\therefore KLMN$ is a cyclic quad [ext \angle of quad = opp int \angle /	✓ R
	<i>buite \angle van vh = teenoorst binne \angle]</i>	
		(1)
OR/OF		
OR/OF		

	$NKL = 180^\circ - K_4$ [adj. suppl.] $\therefore NKL = 180^\circ - NML$ [proved] $\therefore KLMN$ is a cyclic quad [opp. \angle s supplementary]	\checkmark R (1)
11.2.2	In $\Delta LKN \parallel\!\! \Delta KSM$: $\hat{N}_3 = \hat{M}_3 \quad [\angle s \text{ in the same seg} / \angle e \text{ in dieselfde sirkel segm}]$ $\hat{L}_1 = \hat{M}_2 \quad [\angle s \text{ in the same seg} / \angle e \text{ in dieselfde sirkel segm}]$ $= \hat{K}_2 \quad [\text{alt } \angle s; \text{ verw } \angle e; MN \parallel KS]$ $NKL = M\hat{S}K \quad [\angle s \text{ of } \Delta / \angle e \text{ van } \Delta]$ $\Delta LKN \parallel\!\! \Delta KSM$ OR/OF In $\Delta LKN \parallel\!\! \Delta KSM$: $\hat{N}_3 = \hat{M}_3 \quad [\angle s \text{ in the same seg} / \angle e \text{ in dieselfde sirkel segm}]$ $N\hat{K}L = \hat{M}_1 \quad [\text{ext } \angle \text{ of cyclic quad/buite } \angle \text{ van koordevh}]$ $= \hat{S}_2 \quad [\text{corresp } \angle s/\text{ooreenk } \angle e; KS \parallel NM]$ $\Delta LKN \parallel\!\! \Delta KSM \quad [\angle, \angle, \angle]$ OR/OF In $\Delta LKN \parallel\!\! \Delta KSM$: $\hat{N}_3 = \hat{M}_3 \quad [\angle s \text{ in the same seg} / \angle e \text{ in dieselfde sirkel segm}]$ $\hat{K}_4 + N\hat{K}L = \hat{S}_1 + \hat{S}_2 \quad [\angle s \text{ on straight line}/\angle e \text{ op reguitlyn}]$ $\therefore N\hat{K}L = \hat{S}_2 \quad [\hat{K}_4 = \hat{S}_1]$ $\Delta LKN \parallel\!\! \Delta KSM \quad [\angle, \angle, \angle]$	\checkmark S \checkmark R \checkmark S \checkmark S/R \checkmark S (5)
11.2.3	$\frac{LK}{KS} = \frac{KN}{SM} \quad [\Delta LKN \parallel\!\! \Delta KSM]$ $\therefore \frac{12}{KS} = \frac{4}{3}$ $KS = 9 \text{ units}$	\checkmark S \checkmark R \checkmark substitution \checkmark answer (4)
11.2.4	$4SM = 3KN$ $SM = \frac{3(8)}{4}$ $SM = 6$ $\frac{LT}{NL} = \frac{LS}{ML} \quad [\text{line } \parallel \text{ one side of } \Delta / \text{lyn } \parallel \text{ een sy v } \Delta]$ $\frac{LT}{16} = \frac{13}{19}$ $LT = \frac{208}{19} = 10,95$	\checkmark SM = 6 \checkmark S \checkmark R \checkmark answer (4)
		[23]

TOTAL/TOTAAL: 150